

Prevalence of Metabolic Syndrome among Sistanee Ethnic Women

¹Abdoljalal Marjani, ²Najme Shahini, ³Oghuldondi Agh Atabay
and ⁴Robabe Ghiyas Tabari

^{1,3 and 4}Department of Biochemistry and Biophysics, Biochemistry Research Center, Gorgan Faculty of Medicine, Golestan University of Medical Sciences, Gorgan, Golestan province, Iran

²Student Research Committee, Gorgan Faculty of Medicine, Golestan University of Medical Sciences, Gorgan, Golestan province, Iran

Correspondence: Abdoljalal Marjani, Gorgan Faculty of Medicine, Department of Biochemistry and Biophysics, Biochemistry and Metabolic Disorder Research Center, Gorgan Faculty of Medicine, Golestan University of Medical Sciences, Gorgan, Golestan province, Iran
e-mail:abdoljalal@yahoo.com

Abstract

Introduction: The frequency of the metabolic syndrome is increasing throughout the world. The etiology of the metabolic syndrome is dependent on different factors. This study aimed to evaluate the metabolic syndrome among Sistanee ethnic women in Gorgan, North East of Iran.

Methods: The study conducted on the hundred and sixty Sistanee women (20-40years) who were referred to the Health Centers in Gorgan. Metabolic syndrome was diagnosed using Adult Treatment Panel-III (ATP-III) guidelines.

Results: The mean body mass index, waist circumference, systolic blood pressure, diastolic blood pressure, fasting blood glucose and levels were significantly higher in the subjects with metabolic syndrome. The prevalence of low High Density Lipoprotein-cholesterol level, high waist circumference, high triglyceride level, , high blood pressure and high fasting glucose were 23.12%, 22.50%, 16.25%, 12.50% and 0.62%, respectively. Low HDL-cholesterol level (23.12%) and high waist circumference (22.50%) are the most frequent characteristics in comparison to other metabolic components.

Conclusion: The reasonable interpretation for our results is that Sistanee women in this area maybe had lower physical activity. Low HDL-cholesterol and high

waist circumference were the most usual factors of metabolic abnormality among these women.

Keywords: Metabolic syndrome, Sistani ethnic women

Introduction

The metabolic syndrome is described by the clustering of several risk factors for cardiovascular disease (CVD) such as central obesity, glucose intolerance, low level of high-density lipoprotein (HDL), high triglyceride (TG) level, and hypertension (1). There are ethnic differences in the prevalence of metabolic syndrome (2-4). Differences in genetic background, diet, levels of physical activity, age and sex structure all influence the prevalence of both metabolic syndrome and its components (5). The prevalence of metabolic syndrome in adult population worldwide varies from 8 to 24.2% (6-7) in males and from 7 to 46.5% (8-9) in females. Cardiovascular disease is one of the main reasons of death among women in the world (10). The importance of the metabolic syndrome in general populations as a predictor of vascular disease has been confirmed by a number of large prospective epidemiologic studies (11-13). In the United States, the metabolic syndrome has become common (7). The metabolic syndrome is an important public health problem in both developed and developing countries. The prevalence of metabolic syndrome in Europe and among Americans of European ethnic groups change approximately between 20% and 30%, in both gender (2, 5, 7 and 14-15). Some studies show about increasing prevalence of metabolic syndrome in Asia (16). In our area, we do not have enough data on the adult Sistani ethnic group women metabolic syndrome in Gorgan (South East of Caspian Sea), Iran. Therefore, it is very important to set up a study on these women with a risk of metabolic syndrome. The present study aimed to assess the metabolic syndrome among Sistani women in this area.

Methods

This study group included 160 Sistani adult women (Women who speak Sistani language which is native to them) who were referred to the different Health Centers in Gorgan. All the included subjects provided an informed consent. Data were collected by trained interviewers. First of all, a questionnaire was completed at each Health Center by trained interviewers. Demographic information is achieved by a questionnaire. The exclusion criterion was the coexistence of any other serious illness. Exclusion criteria included having hormone replacement therapy, taking drugs such as anti-diabetes and anti-hypertensive anti-lipidemic agents and smokers. A venous blood sample was collected from all the subjects who came after 8-12-hours in the morning after an overnight fast. The samples

were centrifuged for 10 minutes at 3000 rpm. The serum was used for estimating fasting blood glucose, triglycerides, total cholesterol, LDL-cholesterol and HDL-cholesterol concentrations, by biochemical kit using spectrophotometer techniques (Model JENWAY 6105 UV / VIS) in the Biochemistry Research Center (Gorgan Faculty of Medicine). Adult women considered to have metabolic syndrome if they had any three or more of the following, according to the ATP III Criteria: [1]

- A) Abdominal obesity: Waist Circumference >88 cm
- B) Hypertriglyceridaemia: serum triglycerides level > 150 mg/dl
- C) Low HDL-cholesterol: < 50 mg/dl
- D) High blood pressure: SBP > 130 mmHg and/or DBP > 85 mmHg or on treatment for hypertension.
- E) High fasting glucose: serum glucose level > 110 mg/dl or on treatment for diabetes.

Weight was then measured, while subjects were minimally clothed without shoes, using digital scales. Height was measured in standing position using tape meter while the shoulder was in a normal position. Body mass index (BMI) was calculated as weight in kilograms divided by height in meters squared. Those with a BMI of 25.0-29.9 Kg/m² were classified as overweight, whilst those with a BMI \geq 30 Kg/m² were defined as obese. Subjects with BMI greater than 45 Kg/m² were considered very obese (17). Waist circumference was measured at the point halfway between the lower border of ribs and the iliac crest in a horizontal plane (18). Systolic and diastolic blood pressure was measured in sitting position from the right hand. The results were reported as percentages and mean \pm SD. The statistical analysis was done with SPSS- 16 version software. The results were evaluated by using independent student t and Chi square tests. Statistical significance was considered at $P < 0.05$.

Results

The baseline data of the subjects with and without the metabolic syndrome are shown in table 1. The mean body mass index, waist circumference, systolic blood pressure, diastolic blood pressure, fasting blood glucose and levels were significantly higher in the subjects with metabolic syndrome. There were no significant differences in the other parameters in subjects with and without the metabolic syndrome. Prevalence of metabolic syndrome and the components of metabolic syndrome in Sistane ethnic group are shown in table 2. The frequency of metabolic syndrome is shown 23.75%. The prevalence of low High Density Lipoprotein-cholesterol level, high waist circumference, high triglyceride level, , high blood pressure and high fasting glucose were 23.12%, 22.50%, 16.25%, 12.50% and 0.62%, respectively. Low HDL-cholesterol level (23.12%) and high waist circumference (22.50%) are the most frequent characteristics in comparison to other metabolic components. The prevalence of Sistane women with and without metabolic syndrome by age group is shown in table 3. The most age

distribution was in ages between 35-40 years. The highest prevalence of metabolic syndrome was in ages 35-40 years. There was increased frequency of metabolic syndrome from age 20-24 years. The prevalence of metabolic syndrome was significantly high in ages 30-34 years (21.05%) when compared subjects with and without metabolic syndrome. A number of subjects accomplishing the criteria of metabolic syndrome is shown in table 4. Our results showed that 18.12%, 6.87% and 1.87% had three, four and five criteria for metabolic syndrome, respectively.

Table 1 Baseline data of Fars women (Total subjects, subjects with and without metabolic syndrome).

Parameters	Total number of subjects	Subjects with metabolic syndrome	Subjects without metabolic syndrome	P-value
All women, No. (%)	160 (100)	38 (23.75)	122(76.25)	-
Age (years)	30.71±6.80	32.50±6.41	30.11±6.80	0.058
BMI, kg/m ²	26.26±3.98	27.92±4.46	25.76±3.71	0.003
WC, cm	93.51±14.93	100.84±14.17	91.23±14.47	0.001
SBP, mmHg	109.96±19.54	120.76±25.95	106.60±15.74	0.003
DBP, mmHg	67.52±11.98	74.05±13.20	65.49±10.85	0.001
FBS, mg/dl	91.40±26.75	114.35±41.16	84.26±14.34	<0.001
TG, mg/dl	112.44±73.85	159.05±88.21	97.91±62.35	<0.001
T-Chol, mg/dl	187.54±45.54	204.68±70.56	182.20±33.29	0.064
HDL-Chol , mg/dl	40.95±8.94	40.37±6.22	41.13±9.64	0.646
LDL-Chol , mg/dl	126.49±36.92	131.64±55.55	124.88±28.92	0.325

BMI: Body mass index, WC: waist circumference, SBP: systolic blood pressure, DBP: diastolic blood pressure, FBS: fasting blood glucose, TG: triglyceride, T-CHOL: total cholesterol, HDL-CHOL: HDL-cholesterol and LDL-CHOL: LDL-cholesterol

*P value less than 0.05 was considered significant.

Table 2 Prevalence of metabolic syndrome and the components of metabolic syndrome in Sistane women (n=160)

	Num ber	%
metabolic syndrome	38	23.75
Fasting Blood Sugar >110 mg/dl	20	12.50
High Density Lipoprotein-cholesterol < 50 mg/dl	37	23.12
Triglyceride > 150 mg/dl	26	16.25
Waist circumference > 88 cm	36	22.50
Systolic blood pressure >130 mmHg/ Diastolic blood pressure>85 mmHg	1	0.62

Table3. Distribution of Fars women with and without metabolic syndrome by age

Age groups in years	Subjects with MS(n=38)	Subjects without MS(n=122)
20-24 n (%)	3 (7.89)	13 (10.65)
25-29 n (%)	4 (10.52)	17 (13.93)
30-34 n (%)	8 (21.05)*	13(10.65)
35-40 n (%)	23(60.52)	79 (64.75)

MS: Metabolic Syndrome

*P value less than 0.05 was considered significant.

Table 4.Number of subjects accomplishing the criteria of metabolic syndrome

Parameters	subjects(n=160)
3 criteria n (%)	29 (18.12)
4 criteria n (%)	11(6.87)
5 criteria n (%)	3(1.87)
Total criteria n (%)	38(23.75)

Discussion

The frequency of the metabolic syndrome is increasing throughout the world. The etiology of the metabolic syndrome is dependent on different factors such as the high prevalence of abnormal body fat distribution, high triglyceride, insulin

resistance, sociological and environmental, genetic factors and life style. The results of study in Asian people showed these risk factors and its high frequency might start at a young age. The epidemic of metabolic syndrome is worldwide health problem. It is not the only problem of western or Asian countries. Study in Italy showed that a prevalence of metabolic syndrome was 3–3.5% (on the basis of the presence of all five criteria)(20). Studies among Korean and Chinese populations showed that prevalence of metabolic syndrome were 13.8% (21) and 17.8% for females, respectively (22). In Iran, Eshtiaghi et al. showed that prevalence of metabolic syndrome was 18.3% (23). Deilbert (24), Figueiredo Neto (25) is shown that prevalence of metabolic syndrome was 23% and 24%, respectively. In our study the prevalence of the metabolic syndrome is higher than some other studies were done in Italy, Korea, China, Iran ^[20-23], Deilbert (24) and Figueiredo Neto (25), but our study is not in agreement with the studies were done by Ainy (53%) (26) and Heidari et al (44.9%) (27). Studies in Greece and USA have shown that prevalence of metabolic syndrome was similar in both genders (28, 9). Some other studies in Turkey, India, Iran, African Americans, Mexican Americans have shown that women to be much more frequently affected (9, 29), while in France and Australia the metabolic syndrome was found to be more common among men (29). Study on the components of metabolic syndrome showed that the most frequent changes of components of metabolic syndrome was low High Density Lipoprotein-cholesterol (23.12%), which is in agreement with the findings in USA (30), Turkey (31), Italy (32), Canada (33), UK (34) and Iranian population (35-36) that the most common found was high prevalence of low HDL-cholesterol. Some studies have reported that waist circumference is positively associated with the risk of cardiovascular occurrences (37-38). Study of Despres et al showed that extra fat mass rather than excess body weight was highly correlated with abnormal metabolism (39). Our study showed that women with metabolic syndrome had high abdominal obesity. This study showed that waist circumference was elevated among Sistani women with metabolic syndrome. It was been also shown that women were overweight. Changes in central obesity can cause metabolism abnormality and influence health (40). It is important to reduce the risk of cardiovascular disease among these women. It suggests that among women with metabolic syndrome, blood glucose, and blood lipid profile monitoring and changing their life style leading to weight loss by diet and sport (25). Despite lifestyle changes such as an increase in high-fat, high-carbohydrate intake and a decrease in physical activity due to economical alterations in Iran, the metabolic syndrome in women remain an important problem. Study has been shown that 76.3% of females in Iran had physical inactivity (41). Iranian women mostly do less physical activity and overweight and obesity are more common between them (42). The reasonable interpretation for our results is that Sistani women in this area maybe had lower physical activity. Low HDL-cholesterol and high waist circumference were the most usual factors of metabolic abnormality among these women. Prevalence of cardiovascular diseases might be increased. We have shown some related factors

of metabolic syndrome in these women to predict metabolic syndrome in these ethnic groups and help to prevent cardiovascular disease.

Acknowledgments: This study was supported by Golestan University of Medical Sciences. We thank all our colleagues in the Department of Biochemistry, Biochemistry Research Center and Health Centers.

Competing interest: The authors have no conflicts of interest.

Funding: The study was funded by research deputy of Golestan University of Medical Sciences.

References

- [1] Expert panel on detection, evaluation, and treatment of high blood cholesterol in adults. Executive summary of the third report of the national cholesterol education program (NCEP) expert panel on detection, evaluation, and treatment of high blood cholesterol in adults (ATPIII)., *JAMA*, 2001, 285:2486-2497.
- [2] JB Meigs, PW Wilson, DM Nathan, et al, Prevalence and characteristics of the metabolic syndrome in the San Antonio heart and Framingham offspring studies, *Diabetes*, 52(2003), 2160-2167.
- [3]JP Burke, K Williams, SP Gaskill, et al, Rapid rise in the incidence of type 2 diabetes from 1987 to 1996: Results from the San Antonio heart study, *Arch Intern Med*, 41(1999),1450-1456.
- [4] H King, P Zimmet, Trends in the prevalence and incidence of diabetes: Non-insulin dependent diabetes mellitus, *World Health Stat*, 41(1998), 190-196.
- [5] AJ Cameron, JE. Shaw and PZ. Zimmet, The metabolic syndrome: prevalence in worldwide populations, *Endocrinol Metab Clin N Am*, 33 (2004), 351-75.
- [6] A Gupta, R .Gupta, M. Sarna, S. Rastogi, VP. Grupta and K. Kothari, Prevalence of diabetes, impaired fasting glucose and insulin resistance syndrome in an urban Indian population, *Diabetes Res Clin Pract*, 61 (2003), 69-76.
- [7] ES Ford, WH. Giles and WH. Dietz, Prevalence of the metabolic syndrome among US adults: findings from the Third National Health and Nutrition Examination Survey, *JAMA*, 287 (2002), 356-9.
- [8] B Balkau, M. Vernay, L .Mhamdi, M. Novak, D .Arondel and S. Vol, *et al*, The D.E.S.I.R Study Group. The incidence and persistence of the NCEP (National Cholesterol Education Program) metabolic syndrome, The French D.E.S.I.R. study. *Diabetes Metab*, 29 (2003), 526-32.
- [9] A Ramachandran, C .Snehalatha, K. Satyavani, S. Sivasankariand V. Vijay, Metabolic syndrome in urban Asian Indian Adults-a population study using modified ATP III criteria., *Diabetes Res Clin Pract*, 60 (2003), 199-204.
- [10] D Lloyd-Jones , R Adams , M Carnethon, SG De, TB Ferguson, K Flegal, et al, Heart disease and stroke statistics-2009 update: a report from the American Heart Association Statistics Committee and Stroke Statistics Subcommittee, *Circulation*, 119 (2009),480-6.
- [11] J Shepherd, SM. Cobbe, I. Ford, CG. Isles, AR .Lorimerand Mac PW. Farlane, *et al*, Prevention of coronary heart disease with pravastatin in men with hypercholesterolemia, West of Scotland Coronary Prevention Study Group, *N Engl J Med*, 333(1995),1301-7.
- [12] JR Downs, M. Clearfield, S. Weis, E .Whitney, DR. Shapiroand PA. Beerc, *et al*, Primary prevention of acute coronary events with lovastatin in men and women with average cholesterol levels: results of AFCAPS/ TexCAPS. Air Force/Texas Coronary Atherosclerosis Prevention Study, *JAMA*, 279 (1998), 1615-22.
- [13] CM Ballantyne, AG. Olsson, TJ. Cook, MF. Mercuri, TR. Pedersen and J .Kjekshus, Influence of low high-density lipoprotein cholesterol and elevated triglyceride on coronary heart disease events and response to simvastatin therapy in 4S, *Circulation*, 104 (2001), 3046-51.

- [14] Q Qiao, Comparison of different definitions of the metabolic syndrome in relation to cardiovascular mortality in European men and women, *Diabetologia*, 49(2006), 2837-2846.
- [15] B Hildrum, A Mykletun, T Hole , K Midthjell , AADahl, Age-specific prevalence of the metabolic syndrome defined by the International Diabetes Federation and the National Cholesterol Education Program: the Norwegian HUNT 2 study, *BMC Public Health*, 7 (2007), 220.
- [16] JB Meigs, "Invited commentary: insulin resistance syndrome? Syndrome X? Multiple metabolic syndrome? A syndrome at all? Factor analysis reveals patterns in the fabric of correlated metabolic risk factors," *American Journal of Epidemiology*, 10 (2000), 908-912.
- [17] World Health Organization Prevention and Management of the Global Epidemic of Obesity. Report of the WHO Consultation on Obesity. WHO: Geneva, 1998 (Technical Report Series, No. 894).
- [18] M Dalton, A.J.Cameron, P.Z.Zimmet, JE. Shaw, D. D .Jolley, DW Dunstan and TA. Welborn. AusDiab Steering Committee, Waist circumference, waist-hip ratio and body mass index and their correlation with cardiovascular disease risk factors in Australian adults, *J Intern Med*, 254 (2003), 555-63.
- [19] A Misra , NK Vikram, Insulin resistance syndrome (metabolic syndrome) and obesity in Asian Indians: evidence and implications, *Nutrition*, 20 (2004), 482-91.
- [20]. M. Trevisan, J. Liu, F. B. Bahsas, and A. Menotti, "Syndrome X and mortality: a population-based study. Risk Factor and Life Expectancy Research Group," *American Journal of Epidemiology*, 148 (1998), 958-966.
- [21] HM Kim, J Park, SY Ryue, J Kim, The effect of menopause on the metabolic syndrome among Korean women, *The Korean National Health and Nutrition Examination Survey, 2001. Diabetes Care*, 30(2007), 701-6.
- [22] D Gu, K Reynolds, X Wu, et al., "Prevalence of the metabolic syndrome and overweight among adults in China," *The Lancet*, 9468 (2005), 1398-1405.
- [23] R Eshtiaghi, A Esteghamati, M Nakhjavani, Menopause is an independent predictor of metabolic syndrome in Iranian women, *Matutritas*, 65 (2010), 262-266.
- [24] P Deibert, D Konig, MZ Vitolins, U Landmann, I Frey, HP Zahradnik et al, effect of weight loss intervention on anthropometric measures and metabolic risk factors in pre- versus postmenopausal women, *Nutr J*, 6(2007), 31.
- [25] JA Figueiredo Neto, ED Figueredo, ED Barbosa, F Barbosa Fde, GR Costa, VJ Nina, et al, Metabolic syndrome and menopause: Cross-sectional study in Gynecology clinic, *Arq Bras Cardiol*, 95 (2010), 339-45.
- [26] E Ainy, P Mirmiran, S Zahedi Asl, FAzizi, Prevalence of metabolic syndrome during menopausal transition Tehranian women, *Tehran Lipid and Glucose Study (TLGS), Matutritas*, 58(2007), 150-5
- [27] R Heidari, M Sadeghi, M Talaei, K Rabiei, N Mohammadifard, N Sarrafzadegan, Metabolic syndrome in menopausal transition: Isfahan Healthy Heart Program, a population based study, *Diabetol Metab Syndr*, 2(2010), 59.
- [28] VG Athyros, ES Ganotakis, M Bathianaki, I Monedas, IA Goudevenos, AA Papageorgiou et al, MetS-Greece Collaborative Group. Awareness, treatment and control of the metabolic syndrome and its components: a multicentre Greek study, *Hellenic J Cardiol*, 46 (2005), 380-386.

- [29] AJ Cameron, JE Shaw, PZ Zimmet, The metabolic syndrome: prevalence in worldwide populations, *Endocrinology and metabolism clinics of North America. Metabolic Syndrome: Part 1*, 33 (2004), 351-376.
- [30] G Heiss, I Tamir, CE Davis et al, "Lipoprotein-cholesterol distributions in selected North American populations: the lipid research clinics program prevalence study," *Circulation*, 2 (1980), 302-315.
- [31] A Onat, G Surdum-Avci, M Senocak, E Ornek and Y Gozukara, "Plasma lipids and their interrelationship in Turkish adults," *Journal of Epidemiology and Community Health*, 5(1992), 470-476.
- [32] The Research Group ATS-RF2 of the Italian National Research Council, "Distribution of some risk factors for atherosclerosis in nine Italian population samples," *American Journal of Epidemiology*, 113 (1981), 338-346.
- [33] DR MacLean, A. Petrasovits, PW Connelly, M Joffres, B O'Connor and JA Little, "Plasma lipids and lipoprotein reference values, and the prevalence of dyslipoproteinemia in Canadian adults, Canadian Heart Health Surveys Research Group," *Canadian Journal of Cardiology*, 4(1999), 434-444.
- [34] JI Mann, B Lewis, J Shepherd et al, "Blood lipid concentrations and other cardiovascular risk factors: distribution, prevalence, and detection in Britain," *British Medical Journal*, 6638 (1988), 1702-1706.
- [35] F Sharifi, SN Mousavinasab, R Soruri, M Saeini and M Dinmohammadi, "High prevalence of low high-density lipoprotein cholesterol concentrations and other dyslipidemic phenotypes in an Iranian population," *Metabolic Syndrome and Related Disorders*, 3 (2008), 187-195.
- [36] F Azizi, A Esmaillzadeh and P Mirmiran, "Obesity and cardiovascular disease risk factors in Tehran adults: a populationbased study," *Eastern Mediterranean Health Journal*, 6 (2004), 887-897.
- [37] L de Koning, AT Merchant, J Pogue, SS Anand, Waist circumference and waist-to-hip ratio as predictors of cardiovascular events: meta-regression analysis of prospective studies, *Eur Heart J*, 28 (2007), 850-6.
- [38] G Hu, J Tuomilehto, K Silventoinen, C Sarti, S Männistö, P Jousilahti, Body mass index, waist circumference, and waisthip ratio on the risk of total and type-specific stroke, *Arch Intern Med*, 167 (2007), 1420-7.
- [39] JP Despres, I Lemieux, D Prud'homme, Treatment of obesity: need to focus on high risk abdominally obese patients, *BMJ*, 322 (2001), 716-720.
- [40] RA Lobo, Metabolic syndrome after menopause and the role of hormones, *Maturitas*, 60 (2008), 10-18.
- [41] WHO Global infobase, "Iran: Most recent national survey(s) for chronic, non-communicable diseases risk factors," <https://apps.who.int/infobase/report.aspx>.
- [42] P Mirmiran, F Mohammadi, and F Allahveridian Sand Azizi, "Measurement of total energy requirement in adults: prospective Tehran Lipid and Glucose Study, Pajouhandeh," *Journal of Shahid Beheshti University of Medical Sciences*, 6 (2001), 157-166, (Persian).

Received: April, 2012